

COMPLEMENTARY RELATIVITY

Pedro F. González-Díaz.

Instituto de Matemáticas y Física Fundamental
Consejo Superior de Investigaciones Científicas
Serrano 121, 28006 Madrid (SPAIN)

April 7, 1994

There are two possible arenas where events may occur - that of customary space-time and that of a vacuum where events cannot be labelled by space and time, but by a continuum of momentum-energy coordinates. Special theory of relativity has been formulated in a vacuum momentum-energy representation which is equivalent to Einstein special theory of relativity and predicts just the same results as it. Although in this sense such a formulation would be at least classically useless, its consistent extension to noninertial frames produces a momentum-energy metric which behaves as a new dynamical quantity that is here interpreted in terms of a cosmological field. This new field would give rise to interactions transmitted through vacuum, rather than space-time, and is complementary to gravity in that its strength would vary inversely to as that of gravity does. Using a strong-field approximation, we suggest that the existence of these cosmological interactions would induce a shift of luminous energy which could justify the existence of all the assumed invisible matter in the universe, so as the high luminosities found in active galactic nuclei and quasars.

The general framework of theoretical physics may be incomplete in one respect. Whereas quantum mechanics possesses two fully equivalent, though complementary, representations, only the representation that would correspond to position has been worked out in relativity. Although a formulation of special relativity in momentum representation would be fully equivalent to Einstein special relativity in that it predicted exactly the same results, the lack of that representation in general relativity might be hiding some new physics.

In Einstein special relativity, the Minkowskian coordinates t and x have a double function: they serve as labels, but at the same time they also inform us, through Lorentz transformations, about actual time durations and space distances, measurable with clocks and meters. Moreover, although in Einstein relativity momentum and energy can never label real events, they are obtained as actual quantities in its associated relativistic mechanics. Therefore, if we attempted to build up a consistent momentum-energy representation for special relativity, one would expect *no* new predictions from it, but just again the same expressions for time duration and space distances (given in this case as mechanical quantities), now with the coordinates p and e having the double function of serving as labels and informing us about the same actual values of momentum and energy that are predicted by Einstein kinematics. It is in this sense that such a momentum-energy representation of special relativity would be useless.

However, in general relativity this situation could change drastically. For a given representation, we must distinguish here between coordinate labels from proper intervals, entering at the two totally different levels that correspond, respectively, to differential topology and metric geometry. In the usual space-time representation of general relativity, showings of a physical clock are predicted not only by the labels that distinguish events, but also by the metric, and the metric describes at the same time a dynamical quantity - the gravitational field. Likewise, in a consistent momentum-energy representation of general relativity, besides mechanical time durations and space distances, one would expect the emergence of an essentially new quantity - the metric of the momentum-energy continuum, which would help, together the coordinate labels, to construct actual momentum and energy intervals. Such a momentum-energy metric would describe at the same time a new "dynamical" quantity: a field which should be complementary to gravity. It is the physics associated with this new field what may be gained from a complete formulation of relativity.

In what follows I will formulate a momentum-energy represen-

tation for relativity. In order for the resulting full theory to be self-consistent, such a formulation should satisfy the following requirements.

(i) The kinematics of special relativity (i.e. the relations between coordinate labels) in the momentum-energy representation must satisfy all mechanical Einstein four-momentum transformations, and its associated mechanics (i.e. the quantities derived from an action principle) must in turn obey the usual Lorentz transformations.

(ii) Since vacuum is not but what remains after eliminating space-time, we must take it as the arena where events can be continuously labelled by momentum and energy coordinates. Thus, whereas description of a given system in space-time implies that such a system is just a space-time part from a necessarily larger system where at least an external observer is also included, its description in the vacuum momentum-energy continuum requires considering the system as a space-temporal whole where even observers are included, so that no independent evolution of any space-time part of it is possible.

(iii) The nonrelativistic limit $c \rightarrow 0$ of the resulting mechanical relations between time durations and space distances should either produce trivial or known results, or not exist at all. The nonrelativistic limit of the kinematical transformations of momentum and energy must predict values of the vacuum energy which depend on the chosen reference system, and values of the vacuum momentum such that this behaved as an absolute quantity.

The latter requirement needs some further explanation. Consider a system S which evolves uniformly (i.e. at a constant rate $\frac{dp}{de}$) in the vacuum momentum-energy continuum. Since, after requirement (i), its evolution rate is $\frac{dp}{de} = \frac{v}{c^2}$, we can see why the components of momentum must become absolute quantities in the nonrelativistic limit, where energy will still depend on the bare velocity v . For in such a limit, one would not expect the system S with energy e_1 to interact with itself with a different energy e_2 because, then, the maximum rate of signal propagation in momentum-energy, $\frac{1}{c}$, becomes zero. This is precisely what one should expect from the zero-point nonrelativistic oscillator contribution: a nonzero, arbitrary energy and a strictly zero momentum.

Passing to the domain where c is finite, we see that the maximum rate of signal propagation in momentum-energy is no longer zero and, therefore, the momentum components become no longer absolute quantities. This will give rise to the emergence of a purely relativistic interaction of the system S with itself when its energy changes along evolution. We can then introduce momentum-energy

reference systems evolving uniformly relative to each other with relative rates $\frac{v}{c^2}$, so as an extended principle of relativity according to which all the laws of nature are identical in all "inertial" momentum-energy reference systems, if the equations expressing the laws and the events that take place in such reference systems are all described in terms of momenta and energies. Such laws must then be invariant with respect to transformations of momenta and energies from one momentum-energy reference system to another.

A differential interval defined in one of such reference systems can be given by

$$dn^2 = \frac{de^2}{c^2} - dp_x^2 - dp_y^2 - dp_z^2. \quad (1)$$

The principle of relativity for momentum-energy continuum implies that dn will be the same in all inertial momentum-energy systems, and leads to the definition of a proper energy given by

$$de = \frac{de'}{\gamma}, \gamma = (1 - \frac{v^2}{c^2})^{\frac{1}{2}}. \quad (2)$$

Let us consider two inertial momentum-energy reference systems independently evolving with a relative rate $\frac{v}{c^2}$. From the above discussion it follows that if the energy origin is chosen at the point where both systems coincide, and such systems evolve so that their p_x -axes always coincide, then we will have in the limit $c \rightarrow 0$

$$p_x = p'_x, p_y = p'_y, p_z = p'_z, e = e' + p_x v. \quad (3)$$

On the other hand, if c is kept finite, it is easy to see that the transformations that leave invariant the interval are

$$p_x = \frac{p'_x + \frac{v}{c^2}e'}{\gamma}, p_y = p'_y, p_z = p'_z, e = \frac{p'_x v + e'}{\gamma}, \quad (4)$$

which, in turn, coincide with the transformation formulas for momentum energy 4-vector of Einstein relativistic mechanics. Equations (4) lead to expressions for the transformations of velocities, general 4-vectors, and unit 4-velocities, which exactly coincide with those of Einstein relativistic kinematics, and reduce to (3) as $c \rightarrow 0$. Thus, the transformations (4) do satisfy the kinematical parts of the requirements in (i) and (iii).

In order to formulate the relativistic mechanics in momentum-energy representation, let us consider a free system evolving in the momentum-energy continuum. For such a system there should exist a certain integral (the counterpart to action of Einstein relativity) which has the minimum value for actual evolution of the system

in the momentum-energy continuum. This integral must have the form

$$P = -\beta \int_a^b dn = -\frac{\beta}{c} \int_{e_1}^{e_2} de\gamma = \int_{e_1}^{e_2} \tilde{L} de, \quad (5)$$

where $\int_a^b$ is an integral along a momentum-energy world line of the system between two particular events characterizing the momentum of the system when it has energies e_1 and e_2 , and β is some constant that characterizes the system. The coefficient $\tilde{L}$ of de plays the role of a Lagrangian and has the physical dimensions of a time. For P to have the dimensions of an action, unlike Einstein relativity where each system is characterized by its rest energy mc^2 , here each system should be characterized by the complementary quantity to its rest energy, that is its rest time T_0 . We take therefore $\beta = cT_0$, and hence the integral P for a free temporal system becomes

$$P = -T_0 \int_{e_1}^{e_2} de\gamma, \quad (6)$$

with $\tilde{L} = -T_0\gamma$.

Instead of a momentum and an energy, the mechanical system will now be described by a space distance R and a time duration T . Assuming the momentum-energy coordinate space to be homogeneous, so that the properties of the system remain invariant under infinitesimal parallel displacements of rate $\frac{v}{c^2}$ and energy e , the quantities R and T would be conserved and can be obtained using the same Lagrangian principles as in classical mechanics, but in our complementary representation, i.e.

$$R = \frac{\partial \tilde{L}}{\partial(\frac{v}{c^2})} = \frac{T_0 v}{\gamma}, T = R \frac{v}{c^2} - \tilde{L} = \frac{T_0}{\gamma}. \quad (7)$$

We have to check that the relativistic mechanics expressed by (5)-(7) is consistent with the full relativistic picture, i.e. we have to check that by substituting space distance and time duration given in (7), expressed as a 4-vector in terms of the corresponding 4-velocity, in the transformation formulas for a general 4-vector, one obtains usual Lorentz transformations. That this is indeed the case can be readily seen by using the principle of least action, $\delta P = 0$, and $dn = (dp_i dp^i)^{\frac{1}{2}}$, with $p^0 = \frac{e}{c}$, $p^1 = p_x$, $p^2 = p_y$, $p^3 = p_z$. We then obtain $\delta P = -T_0 s_i \delta p^i$, where $s_i = \frac{dp^i}{dn} = u_i$, u_i being the Einstein unit 4-velocity. It follows that

$$x_i = -\frac{\partial P}{\partial p^i} = (cT, R) = T_0 u_i \quad (8)$$

is the distance 4-vector. It turns out that the square of the length of momentum 4-vector, $(p^0)^2 - \sum_{\alpha=1}^3 (p^\alpha)^2$, is invariant under transformations (4). Generalizing to any 4-vector A^i which transforms like

the components of the momentum 4-vector under (4), we recover the usual transformation formulas for 4-vectors of Einstein relativity. It is now immediately seen that by substituting (8) into such formulas one obtains usual Lorentz transformations. This completes fulfillment of requirement (i).

We also note that the formula for T in (7) has no general nonrelativistic counterpart. In fact, in the limit $c \rightarrow \infty$, we obtain from (7)

$$T \approx T_0 + \frac{T_0 v^2}{2c^2} \approx T_0, R \approx T_0 v, \quad (9)$$

i.e. the nonrelativistic limit of T and R reduces, respectively, to the rest time and a distance-velocity law which may be trivially interpreted as the customary definition of velocity. Rest time T_0 would correspond to the period of the wave that quantum theory associates with the corpuscular characteristics of the system.

On the other hand, it follows from (7)

$$R = T v \quad (10)$$

$$T^2 c^2 = R^2 + T_0^2 c^2. \quad (11)$$

Expression (9) should now correspond to the relativistic expression for the definition of velocity. We finally note that (11) must correspond to the analogue of the usual relativistic Hamiltonian in our complementary formalism for relativity. If we express time T in terms of the distances R , then we have a complementary relativistic Hamiltonian

$$T \equiv H_T = \frac{1}{c} (R^2 + T_0^2 c^2)^{\frac{1}{2}}, \quad (12)$$

which has the physical dimension of a time. Law (12) must correspond to the Minkowskian function F which is the conjugate counterpart to Hamiltonian and whose existence has been recently suggested [1].

We still have to check that our mechanical relation (11) satisfies requirement (iii). In the limit $c \rightarrow \infty$ (11) again gives rise to no nonrelativistic complementary Hamiltonian, except for the rest time of the system. We conclude therefore that law (12) cannot have any mechanical effects in our familiar low-velocity physics, other than making it possible the existence of the quantum-mechanical wave-particle dual character for the given system. When applied to the whole universe, this law will describe the cosmological evolution in the vacuum momentum-energy continuum. A quantum-mechanical wave equation can also be derived from (11) by introducing the operators $\hat{T} = i\hbar \frac{\delta}{\delta e}$, $\hat{R} = i\hbar \frac{\delta}{\delta p}$. Using a wave function $\Upsilon \equiv \Upsilon(p, e)$, we

obtain

$$-\hbar^2 \frac{\partial^2 \Upsilon}{\partial e^2} = \frac{1}{c^2} (-\hbar^2 \frac{\partial^2}{\partial p^2} + T_0^2 c^2 + V(p)) \Upsilon, \quad (13)$$

where we have introduced a generic potential $V(p)$. This is the complementary-relativity analogue to the Klein-Gordon equation. If, as it is the case for the whole universe, the system is closed, then one would expect a discrete T -spectrum which would associate with an infinite set of universes frozen at the given values of T . This spectrum would only become continuous in the classical region that corresponded to very large values of T . We finally note that the quantum description of systems that show time asymmetry could only be accounted for whenever we assume a half-integer intrinsic angular momentum for the whole system, so that, instead of (13), one would have a Dirac-like wave equation

$$(\gamma^i \frac{\partial}{\partial p^i} + cT_0 + V(p)) \Upsilon(p) = 0, \quad (14)$$

with γ^i the 4×4 Dirac matrices, which is invariant under $e \rightarrow -e$, but not under $T \rightarrow -T$. Indeed, just as for antimatter in momentum representation, the negative time states could not be physically ignored, since there is nothing to prevent a system from making a transition from a state of positive time to a state of negative time.

Extending the above formalism to noninertial momentum-energy reference systems on which general covariance is imposed [2], we are left with an invariant momentum interval

$$dn = f_{ij} dp^i dp^j, \quad (15)$$

where, analogously to metric tensor g_{ij} with respect to space-time and gravitational field, relative to an arbitrarily chosen system of momentum-energy coordinates, the functions f_{ij} describe the metrical relations of the vacuum momentum-energy continuum and also a field which is complementary to gravity. Such a field is what we have gained from formulating relativity in momentum-energy representation. But, what does this new field represent?. First of all, according to our requirement in (ii), it must govern either self-interaction of the wholeness of independent space-temporal systems evolving in vacuum, rather than space-time, or the interaction of two or more of such systems. Moreover, if we consistently extend all our above requirements to also embrace noninertial reference systems, so that it is only this new field that constitutes all of the new physics gained, then the whole mathematics of general relativity would exactly re-appear for noninertial momentum-energy relativity by simply replacing g_{ij} for f_{ij} and x^i for p^i within the realm of Riemmanian geometry, and finally we shall arrive to obtain field

equations exactly like Einstein equations, but expressed in terms of f_{ij} , p^i and the *space-time* 4-tensor, rather than the usual g_{ij} , x^i and *momentum-energy* 4-tensor (see Eqn. (18)). These equations can then be interpreted as requiring that the space-temporal characteristics of a system evolving in vacuum cause momentum-energy curvature C , or alternatively, that such characteristics of the system evolve in a certain way in response to the existence of curvature C . Consider now two of such systems, each causing a certain amount of vacuum curvature, and evolving in a certain way through momentum-energy in response to the C -curvature caused by the other. The net result would be that each system influences the other in such a way that their mutual vacuum momentum separation will always tend to decrease. It is this influence that describes the complementary field. Moreover, if the mutual influence is very small (large), then the vacuum momentum separation between the two systems will become huge (tiny) and, by the uncertainty principle, their mutual space distance very small (large). Hence, the stronger gravity, the weaker its complementary field, and *vice versa*. It follows that the new field would manifest as a repulsive interaction in space-time. It appears then that the new interaction is interpretable as a *cosmological* field which influences every possible system evolving in vacuum momentum-energy continuum. The ultimate evolution of a given system will be the result of the competition between the gravitational and cosmological interactions to which it is subject.

Of most interest is the expression which describes the evolution of a system in a cosmological field with potentials f_{ij} , i.e.

$$\frac{d^2 p^i}{dn^2} + \Upsilon_{kl}^i u^k u^l = 0, \quad (16)$$

with Υ_{kl}^i the complementary affine connection in momentum-energy

$$\Upsilon_{kl}^i = \frac{1}{2} f^{im} \left(\frac{\partial f_{mk}}{\partial p^l} + \frac{\partial f_{ml}}{\partial p^k} - \frac{\partial f_{kl}}{\partial p^m} \right) \quad (17)$$

and the u^i are the conventional 4-velocities in space-time which equal their counterparts s^i for momentum-energy. We shall denote the momentum-energy itinerary followed by the system as a *cosmodesic*.

By a suitable choice of the momentum system we can always make all $\Upsilon_{kl}^i = 0$ at an arbitrary value of momentum-energy, and this would mean the elimination of the cosmological field in the given infinitesimal element of momentum-energy. At such an element, it would happen that the gravitational field becomes infinite, and *vice versa*, at the space-time points where the gravitational field

is made to vanish the cosmological field would diverge. This discriminated singularities of two essentially different kinds: spacetime singularities at which gravity becomes infinite and the cosmological field vanishes, and momentum-energy singularities at which gravity vanishes and the cosmological field diverges.

On the other hand, one would have as cosmological field equations

$$C_{ik} = \frac{4\pi K}{3}(S_{ik} - \frac{1}{2}f_{ik}S), \quad (18)$$

where C_{ik} is the Ricci curvature tensor as expressed in terms of f_{ik} and p^i , instead of g_{ik} and x^i , respectively, K is a cosmological coupling constant having the dimension of a conventional force, and S is a *space-time* 4-tensor. A more detailed study of the cosmological field and its relation with gravitation will be considered elsewhere. Here, we shall only regard the cosmological field when all involved velocities are small compared with the velocity of light. In such a case, the components of S_i^k are given by

$$S_i^k = \tau_0 u_i u^k, \quad (19)$$

with τ_0 the time momentum-density of the system, i.e. the time that characterizes the system in a unit momentum-volume, $\tau_0 = \frac{T_0}{V_p} = \frac{3T_0}{4\pi p^3}$. Let us introduce then a nonrelativistic cosmological field and denote it by ρ . In general, such a field will depend on coordinates p and e and have the dimension of the inverse of a squared velocity. The nonrelativistic complementary Lagrangian with dimension of a time in the presence of a cosmological field will be given by

$$\tilde{L} \approx T_0(\frac{v^2}{2c^2} - c^2\rho - 1) \quad (20)$$

and is associated with a complementary nonrelativistic action

$$P \approx -T_0 c \int de (\frac{1}{c} + c\rho - \frac{v^2}{2c^3}). \quad (21)$$

It follows that in this limiting case the components of the complementary metric tensor become

$$f_{00} = c^2(c^2\rho^2 + 2\rho), f_{\alpha\alpha} = 1 + c^2\rho, \alpha = 1, 2, 3 \quad (22)$$

In the limit $\frac{v}{c} \rightarrow 0$, the components of 4-velocity become $u^\alpha = 0$, $u^0 = u_0 = 1$, so that the only surviving component of tensor S_i^k is $S_0^0 = \tau_0$. From (18) we then obtain

$$C_0^0 = \frac{4\pi K}{3}\tau_0.$$

We note that the terms containing derivatives of the affine connections Υ_{kl}^i with respect to $\frac{e}{c}$ involve extra power c and, unlike for Einstein gravity, are large as compared to the derivative with respect to momenta p^α . Therefore the term that dominates in the nonrelativistic limit the Ricci curvature tensor

$$C_0^0 = \frac{\partial \Upsilon_{00}^\alpha}{\partial p^\alpha} - \frac{\partial \Upsilon_{0\alpha}^\alpha}{\partial p^0} + \Upsilon_{00}^\alpha \Upsilon_{\alpha\beta}^\beta - \Upsilon_{0\alpha}^\beta \Upsilon_{0\beta}^\alpha \quad (23)$$

is the negative second term on the r.h.s. It follows then that in this limit $C_0^0 \approx -\frac{\partial \Upsilon_{0\alpha}^\alpha}{\partial p^0}$. Now, unlike for the gravitational field [3], one should expect that the assumption of small velocities implies the requirement that the cosmological field be strong, for if this were not the case an individual subsystem located in it would acquire a high arbitrary velocity with respect to the whole system, which is contrary to the very notion of the cosmological field whose effect would instead be slowing all individual independent subsystem motions to produce an overall joint evolution pattern. Therefore, one expected that when $v \rightarrow c$ the cosmological field would vanish, and for small velocities $c^2\rho \gg 1$. Hence

$$\Upsilon_{0\alpha}^\alpha = \frac{1}{2} f^{\alpha\beta} \frac{\partial f_{\alpha\beta}}{\partial p^0} \approx -\frac{1}{2} c \frac{\dot{\rho}}{\rho}, \quad (24)$$

where $\dot{\rho} = \frac{d\rho}{de}$. We obtain finally

$$\frac{d}{de} \left(\frac{\dot{\rho}}{\rho} \right) \approx \frac{8\pi K \tau_0}{3c^2}. \quad (25)$$

If the system is characterized by a single time T_0 , then we obtain an asymptotic expression for the cosmological field

$$\rho \approx \rho_0 \exp\left(\frac{KT_0 c^2}{P v^2}\right), \quad (26)$$

where we have used $e^2 = \frac{P^2 c^4}{v^2}$, and ρ_0 is an integration constant. Potential (26) would, as expected, correspond to a repulsive field. Note that the larger the "momentum separation", the smaller the value of the potential, and the larger the absolute value of the involved gravitational potential.

Let us consider now the case of a constant cosmological field, meaning by that a cosmological field which does not depend on vacuum energy coordinate p^0 : $f_{ik} \neq f_{ik}(p^0)$. In this case the coordinate p^0 should be a true *vacuum energy* relating to the complementary proper energy e by $e = c f_{00}^{\frac{1}{2}} p^0$. In the case of a strong cosmological field this relation would reduce to

$$e \approx c p^0 (1 + c^2 \rho) \quad (27)$$

for which $c^2\rho \gg 1$ and the first of expressions (22) have been used.

Let us look then at the momentum-energy propagation in a large constant cosmological field of a light ray with an *energy frequency* Ω [4] which would generally be given by the derivative with respect to energy coordinate of the phase-eikonal in momentum-energy continuum $\varphi = -r_i p^i + \beta$ (where β is an arbitrary constant) for a momentum-energy "plane wave" $g = g_0 e^{i\varphi}$ which corresponded to just one space-time point. As expressed in terms of the vacuum energy p^0 , the energy frequency becomes $c\Omega_0 = -\frac{\partial\varphi}{\partial p^0}$, and if we express it in terms of the proper energy e , then we have

$$c\Omega = -c\frac{\partial\varphi}{\partial e} = -f_{00}^{-\frac{1}{2}}\frac{\partial\varphi}{\partial p^0} \approx \frac{\Omega_0(1 - \frac{1}{c^2\rho})}{c\rho}. \quad (28)$$

Thus, if a ray of light is emitted at a point where the cosmological potential is ρ_1 and the energy frequency is Ω , then upon arriving at a point where the potential is ρ_2 it will have an approximate energy frequency $\frac{\Omega\rho_1}{\rho_2}$. For an observer at the arriving point the energy frequency would then be shifted by an amount

$$\Delta\Omega \approx \Omega\left(\frac{\rho_1}{\rho_2} - 1\right), \quad (29)$$

which corresponded to a cosmological proper-energy shift given by

$$\Delta E \approx \left(\frac{\rho_2}{\rho_1} - 1\right)E, \quad (30)$$

where E is the proper energy at the emission point where the cosmological potential is ρ_1 .

In order to analyse the possible observational meaning of (30), let us consider the light coming to our galaxy from another luminous galaxy. It appears that the light emitted from inner regions of the latter galaxy should be produced in a physical environment where the local gravitational velocities would generally be comparable to those occurring in our own; therefore, for such a light, both the gravitational and cosmological fields would be expected to be of the same intensity for the two galaxies, and $\Delta E \approx 0$. However, as the light source separates from the core and enters the outer regions of the emitting galaxy, its gravitational atmosphere became weaker and hence its related velocities would be smaller. Thus, the cosmological field at the outskirts of the emitting galaxy should be expected to be stronger than at the receiving point on our galaxy. One then would generally expect $\Delta E < 0$ for light coming from galactic halos.

An observer in our galaxy who were unaware of (30) would therefore interpret the amount of luminous matter in the emitting galaxy

to become smaller than it actually is as the source is shifted from galactic centers first towards the halos, and then deep into the intergalactic media where the cosmological field became strongest. It appears that, rather than interpreting this indeed observed effect as caused by the existence of more or less exotic dark matter [5], it should be attributed to the cosmological shift (30) which originates from the noninvariance of the proper energy under propagation in momentum-energy continuum.

On the other hand, one would not exclude the observational emergence of just the opposite effect; i.e. the effect arising from physical situations where the cosmological field at the source is weaker than in our galaxy. Such situations would in principle appear when the emitting regions possessed very strong gravitational fields and correspond to positive cosmological shifts, $\Delta E > 0$. An observer in our galaxy would then attribute an anormally high luminosity to the source, typically perhaps an active galactic nucleus or a quasar. Of course, all the above discussion is rather qualitative as the approximate expression for the cosmological energy shift given by (30) only is strictly valid for cosmological fields sufficiently strong as for $c^2\rho$ to still be much greater than unity.

Extrapolating to the point where all velocities vanish, a true momentum-energy singularity would be reached where $\rho \rightarrow \infty$ and $\Delta E = -E$. So, no luminous matter could be detected from that point by any observers at any regions. If we assumed this to be big bang singularity, then immediately after it the huge value of the repulsive potential ρ given by (26) would drive the universe through a very short-lived phase of exponential expansion which rapidly damped off to smoothly enter a regime where the cosmological potential is small enough to allow dominance of Einstein gravitational field, and hence of an evolution that proceeds in accordance with conventional Friedmann expansion. All memory of the cosmological field left on the universe as a whole would then be slowly erased as expansion goes on, and correspond to the usual cosmological term Λ .

Acknowledgements. The author is grateful to *CAICYT* for support under Research Project N PB91-0052. I also thank Carmen Siguenza for her useful comments, constructive criticisms and constant encouragment, and José María Gómez Soto for his interest and deep understanding.

References

- 1 P.F. González-Díaz, Phys. Lett. B317 (1993) 36.
- 2 A. Einstein, *The Meaning of Relativity* (Chapman and Hall, Bristol, 1967); in: *The Principle of Relativity* (Dover, New York, 1952).
- 3 L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press, Oxford, 1975).
- 4 What plays the role of a space-time wave in momentum-energy should be a phenomenon repeating itself along the energy coordinate, with energy frequency Ω . Light signals propagated in momentum-energy at the maximum rate $\frac{1}{c}$, and manifested either as waves with frequency Ω , or as discrete *time quanta* where time concentrated so that $T = h\Omega$. These definitions are necessary for a consistent quantization of systems in momentum-energy, since only they may allow that a periodic phenomenon occurring in momentum-energy satisfies the counterpart for complementary relativity of the de Broglie's theorem of phase harmony [L. de Broglie, *Recherches sur la Theorie des Quanta* (Masson and Cie., Paris, 1963)]. Clearly, by using these definitions, frequency $\Omega_1 = \frac{T_0\gamma}{h}$ (from (4)) relative to a fixed observer will always keep itself in phase ($\varphi = \frac{T_0\gamma e}{h}$) with a wave of energy frequency $\Omega = \frac{T_0}{h\gamma}$ (from (7)), for such an observer. The latter wave propagated at a rate $\frac{1}{v} > \frac{1}{c}$ and carried only phase. It can be also shown that the group rate for such phase waves is $\frac{v}{c^2}$, i.e. just the propagation rate of the system in momentum-energy.
- 5 P.J.E. Peebles, *Principles of Physical Cosmology* (Princeton University Press, Princeton, 1993).